

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## SDCBP monoclonal antibody (M01), clone 2C12

Catalog Number: H00006386-M01

Regulation Status: For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against a partial recombinant SDCBP.

Clone Name: 2C12

 $\label{eq:mmunogen: SDCBP (NP_005616, 1 a.a. $\sim$ 100 a.a)} partial recombinant protein with GST tag. MW of the$ 

GST tag alone is 26 KDa.

## Sequence:

MSLYPSLEDLKVDKVIQAQTAFSANPANPAILSEASAPI PHDGNLYPRLYPELSQYMGLSLNEEEIRANVAVVSGA PLQGOLVARPSSINYMVAPVTGND

Host: Mouse

Reactivity: Human

Applications: ELISA, IF, IHC-P, IP, S-ELISA, WB-Ce,

WB-Re, WB-Tr

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 6386

Gene Symbol: SDCBP

Gene Alias: MDA-9, ST1, SYCL, TACIP18

**Gene Summary:** The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The

syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq]

## References:

1. Novel Role of MDA-9/Syntenin in Regulating
Urothelial Cell Proliferation by Modulating EGFR
Signaling. Dasgupta S, Menezes ME, Das SK, Emdad L,
Janjic A, Bhatia S, Mukhopadhyay ND, Shao C, Sarkar
D, Fisher PB Clin Cancer Res. 2013 Sep
1;19(17):4621-33. doi:
10.1158/1078-0432.CCR-13-0585. Epub 2013 Jul 19.

2. MDA-9/Syntenin and IGFBP-2 Promote Angiogenesis in Human Melanoma. Das SK, Bhutia SK, Azab B, Kegelman TP, Peachy L, Santhekadur PK, Dasgupta S, Dash R, Dent P, Grant S, Emdad L, Pellecchia M, Sarkar D, Fisher PB. Cancer Res. 2013 Jan 15;73(2):844-54. doi: 10.1158/0008-5472.CAN-12-1681. Epub 2012 Dec 10.

3. Raf kinase inhibitor RKIP inhibits

MDA-9/syntenin-mediated metastasis in melanoma. Das SK, Bhutia SK, Sokhi UK, Azab B, Su ZZ, Boukerche H, Anwar T, Moen EL, Chatterjee D, Pellecchia M, Sarkar D, Fisher PB. Cancer Res. 2012 Oct 11. [Epub ahead of print]